

An Overlooked Electrocardiogram Pattern: de Winter Pattern

Ömer IŞIK^a, Ali Han ÖZKAN^b

^aFırat University Faculty of Medicine, Department of Cardiology, Elazığ, Türkiye

^bTokat Gaziosmanpaşa University Faculty of Medicine, Tokat, Türkiye

ABSTRACT Electrocardiogram (ECG) is an imaging tool used in the diagnosis of acute coronary syndrome. ST elevation is seen in total occlusion of the coronary artery. However, there are a number of high-risk ECG patterns that do not meet the criteria for ST-segment elevation myocardial infarction (STEMI) but are still suggestive of coronary artery occlusion. An ECG pattern called the de Winter pattern is an ECG pattern that indicates severe occlusion of the proximal left anterior descending artery. In the de Winter pattern, there is an upward sloping ST segment at the J point that continues into long, positive symmetric T waves in the precordial leads. This ECG pattern is equivalent to a STEMI; therefore these patients should be triaged for emergency intervention. This case report describes a patient with this finding and reviews the current literature.

Keywords: Myocardial infarction; de Winter pattern; acute coronary syndrome; coronary artery disease

Ischemic heart disease can be treated across a broad clinical spectrum, from silent ischemia to sudden death. Most deaths are related to acute coronary syndromes (ACS) and their complications.¹ Although ACS are a condition in which there are serious advances in treatment, they continue to be very important in terms of patient potential. Rapid diagnosis and intervention are extremely important for adequate treatment.^{2,3} Although the differential diagnosis of ACS continues to be important, studies conducted over the years have shown that there are different acute coronary conditions and have revealed new electrocardiographic patterns.^{4,5} One of these patterns is the de Winter pattern, first published by de Winter et al. in 2008, which includes specific changes in electrocardiography (ECG) and suggests that left anterior descending (LAD) artery occlusion should be treated in the acute phase. As suggested in the article by de Winter et al., this includes at least 1 mm nega-

tivity of the ST segment starting from the J point in the precordial leads, pointed and symmetrical T waves, and small ST elevation (0.5 mm-1 mm) in aVR.^{6,7} de Winter pattern occurs as an acute manifestation of coronary artery occlusion, such as ST-segment elevation myocardial infarction (STEMI). Although LAD occlusion is predominantly seen, there are also studies involving other coronary arteries.^{8,9} de Winter pattern is a rare condition and has been observed to occur in patients with anterior myocardial infarction at rates ranging between 2% and 3.4% in different case series.⁹

In this case, we presented a case with de Winter pattern in the ECG.

CASE REPORT

A 31-years-old male patient presents to the emergency department with a complaint of compressive

TO CITE THIS ARTICLE:

Işık Ö, Özkan AH. An overlooked electrocardiogram pattern: de Winter pattern. Türkiye Klinikleri J Case Rep. 2024;32(3):77-80.

Correspondence: Ömer IŞIK

Fırat University Faculty of Medicine, Department of Cardiology, Elazığ, Türkiye

E-mail: drr.omr@gmail.com

Peer review under responsibility of Türkiye Klinikleri Journal of Case Reports.

Received: 08 Feb 2024

Accepted: 14 Jun 2024

Available online: 24 Jun 2024

2147-9291 / Copyright © 2024 by Türkiye Klinikleri. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).






FIGURE 1: Electrocardiogram of the patient at the time of admission to the emergency department.

chest pain that has been present for two days. In the patient's family history, it was learned that his parents were cousins and both had a history of ACS under the age of 40. The patient has a 5 pack/year smoking history. Our patient has no known history of additional diseases. In the 12-lead ECG taken at our patient's admission to the emergency department, there were high-protruding symmetrical T waves in the precordial leads, >1 mm upward-sloping ST segment depression at the J point, absence of ST segment elevation, and ST segment elevation (0.5 mm-1 mm) in aVR (Figure 1). In the echocardiography (ECHO), the anterior region was hypokinetic and the ejection fraction was found to be 35%. When the patient's current clinical condition and ECG and ECHO findings were evaluated, the patient was taken to the coronary angiography (CAG) laboratory with the diagnosis of ACS. In the CAG performed on the patient, a 90% thrombosed lesion in the LAD was followed by a 30% lesion in the right coronary artery (RCA) and a plaque in the circumflex artery (Figure 2, Figure 3, Figure 4). A stent was applied to the patient's responsible lesion in the LAD. The procedure was completed successfully without complications. In the patient's 12-lead ECG taken after CAG, the ST segment in aVR in the precordial segments returned to the isoelectric line and the ST segment elevation in aVR returned to the isoelectric line (Figure 5).




FIGURE 2: Angiographic image of the patient's right coronary artery vessel.

Informed consent form was obtained from the patient.

DISCUSSION

The purpose of this case report is to emphasize that this ECG pattern is as important as STEMI and that these patients should undergo immediate percutaneous coronary intervention because the de Winter pattern occurs as a consequence of significant narrowing of the proximal LAD. The de Winter pattern




FIGURE 3: Angiographic image of the responsible lesion in the patient's left anterior descending vessel.




FIGURE 4: Angiography of the left anterior descending vessel after the procedure.




FIGURE 5: Electrocardiogram of the patient after the procedure.

was identified in 2% of patients with myocardial infarction in the LAD and was associated with a mortality of 27% within the first week.⁹ Initially, it was described as a pattern obtained on average 1.5 hours after the onset of chest pain. Later, it was suggested that the de Winter pattern was on a continuum with ECG changes of subendocardial ischemia and transmural infarction of STEMI.¹⁰ Ghaffari et al. reported RCA occlusion in a patient with de Winter pattern in inferior leads.¹¹ Chen et al. reported a case of de Winter ECG pattern in a patient who underwent primary percutaneous coronary intervention to the RCA and perioperatively developed new-onset chest pain and de Winter complexes, staging PTCA to the LAD and Cx.¹² The resulting coronary angiographies did not

reveal any significant restenosis in the epicardial coronaries or stent thrombosis, and the development of such de Winter complexes has been attributed to perioperative microvascular ischemia due to increased cardiac biomarkers.

The mechanism of the de Winter model has not yet been determined, but it has been suggested that it may develop due to lack of activation of potassium-sensitive ATPase channels due to ischemic exhaustion.¹² In the hyperacute phase of STEMI, tall, symmetrical T waves are caused by subendocardial ischaemia. In the acute phase of STEMI, ST-segment elevation is caused by transmural ischaemia. However, the sensitivity to ischaemia is different between endocardium and epicardium, es-

pecially the cells at the junction between the intermediate myocardium and the endocardium. Hypoxia in the mid-myocardium may be the main cause of ECG manifestation of the de Winter pattern.¹³

de Winter patterns are high-risk ECG patterns associated with a large area of transmural infarction, necessitating early diagnosis and rapid intervention by the healthcare team. Primary PTCA reperfusion should be the mainstay of therapy. de Winter T waves should be clearly articulated in training courses and guidelines so that patients can receive appropriate treatment in a timely manner, reducing morbidity and mortality.

Source of Finance

During this study, no financial or spiritual support was received neither from any pharmaceutical company that has a direct connection with the research subject, nor from a company that provides or produces medical instruments and materials which may negatively affect the evaluation process of this study.

Conflict of Interest

No conflicts of interest between the authors and / or family members of the scientific and medical committee members or members of the potential conflicts of interest, counseling, expertise, working conditions, share holding and similar situations in any firm.

Authorship Contributions

All authors contributed equally while this study preparing.

REFERENCES

1. Timmis A, Townsend N, Gale CP, Torbica A, Lettino M, Petersen SE, et al; European Society of Cardiology. European Society of Cardiology: Cardiovascular Disease Statistics 2019. *Eur Heart J*. 2020;41(1):12-85. Erratum in: *Eur Heart J*. 2020;41(47):4507. [[Crossref](#)] [[PubMed](#)]
2. Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno H, et al; ESC Scientific Document Group. 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: The Task Force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J*. 2018;39(2):119-77. [[PubMed](#)]
3. O'Gara PT, Kushner FG, Ascheim DD, Casey DE Jr, Chung MK, de Lemos JA, et al. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2013;61(4):e78-e140. [[PubMed](#)]
4. Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, Bax JJ, Morrow DA, et al; Executive Group on behalf of the Joint European Society of Cardiology (ESC)/American College of Cardiology (ACC)/American Heart Association (AHA)/World Heart Federation (WHF) Task Force for the Universal Definition of Myocardial Infarction. Fourth Universal Definition of Myocardial Infarction (2018). *J Am Coll Cardiol*. 2018;72(18):2231-64. [[Crossref](#)] [[PubMed](#)]
5. Vilela EM, Sampaio F, Dias T, Barbosa AR, Primo J, Caeiro D, et al. A critical electrocardiographic pattern in the age of cardiac biomarkers. *Ann Transl Med*. 2018;6(7):133. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
6. de Winter RJ, Verouden NJ, Wellens HJ, Wilde AA; Interventional Cardiology Group of the Academic Medical Center. A new ECG sign of proximal LAD occlusion. *N Engl J Med*. 2008;359(19):2071-3. [[Crossref](#)] [[PubMed](#)]
7. Vilela EM, Caeiro D, Primo J, Braga P. A pivotal electrocardiographic presentation: reading between the lines. *Neth J Med*. 2019;77(8):297. [[PubMed](#)]
8. Morris NP, Body R. The De Winter ECG pattern: morphology and accuracy for diagnosing acute coronary occlusion: systematic review. *Eur J Emerg Med*. 2017;24(4):236-42. [[Crossref](#)] [[PubMed](#)]
9. de Winter RW, Adams R, Amoroso G, Appelman Y, Ten Brinke L, Huybrechts B, et al. Prevalence of junctional ST-depression with tall symmetrical T-waves in a pre-hospital field triage system for STEMI patients. *J Electrocardiol*. 2019;52:1-5. [[Crossref](#)] [[PubMed](#)]
10. Goebel M, Bledsoe J, Orford JL, Mattu A, Brady WJ. A new ST-segment elevation myocardial infarction equivalent pattern? Prominent T wave and J-point depression in the precordial leads associated with ST-segment elevation in lead aVr. *Am J Emerg Med*. 2014;32(3):287.e5-8. [[Crossref](#)] [[PubMed](#)]
11. Ghaffari S, Pourafkari L, Nader ND. "de Winter" electrocardiogram pattern in inferior leads in proximal right coronary artery occlusion. *Arch Cardiol Mex*. 2021;91(3):366-8. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
12. Chen S, Wang H, Huang L. The presence of De Winter electrocardiogram pattern following elective percutaneous coronary intervention in a patient without coronary artery occlusion: a case report. *Medicine (Baltimore)*. 2020;99(5):e18656. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
13. Xu J, Wang A, Liu L, Chen Z. The de winter electrocardiogram pattern is a transient electrocardiographic phenomenon that presents at the early stage of ST-segment elevation myocardial infarction. *Clin Cardiol*. 2018;41(9):1177-84. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]