

# Mycotoxin Analysis in Pollen and Bee Bread Samples: Experimental Study

## Polen ve Arı Ekmeği Örneklerinde Mikotoksin Analizi: Deneysel Araştırma

<sup>id</sup> Sibel SİLİCİ<sup>a</sup>, <sup>id</sup> Uğur KESKİN<sup>a</sup>

<sup>a</sup>Erciyes University Faculty of Agriculture, Department of Agricultural Biotechnology, Kayseri, Türkiye

**ABSTRACT Objective:** Apitherapy is the use of bee products such as honey, pollen, bee bread, royal jelly, propolis and bee venom to protect against diseases and support healing. However, it is important that these products do not contain residues and contaminants. For this reason, this study aimed to determine the contaminant content by performing aflatoxin and ochratoxin analysis on pollen and bee bread samples obtained from different geographical regions. **Material and Methods:** In this study, moisture, dry matter, ash, protein and mycotoxin content of 18 pollen (Kırşehir, Nevşehir, Kayseri, Aydın, Silifke, Niğde, Bursa, Denizli, Isparta, Kütahya, İzmir, Afyon) and 2 bee bread samples (Kırşehir and Nevşehir) obtained from beekeepers were analyzed. **Results:** The moisture content of the samples is between 13.82% and 29.10% (average 17.80), dry matter content is between 70.90 and 86.18 (average 82.20), ash is between 1.60% and 3.32% (average 2.11) and protein content is between 15.06% and 27.95% (average of 20.49). Aflatoxin B1, Aflatoxin B2, Aflatoxin G1, Aflatoxin G2 and total Aflatoxin (B1+B2+G1+G2) levels were not detected above the detection limit (0.60 µg/kg) in any of the samples tested. **Conclusion:** According to the results obtained, the mycotoxin analysis values of pollen and bee bread samples obtained from different regions of our country were found to be in accordance with the Turkish Food Codex Limits.

**Keywords:** Pollen; bee bread; chemical analysis; aflatoxin; mycotoxin

**ÖZET Amaç:** Apiterapi bal, polen, arı ekmeği, arı sütü, propolis ve arı zehiri gibi arı ürünlerinin hastalıklardan koruma ve tedaviye destek amacıyla kullanımınıdır. Ancak bu ürünlerin kalıntı ve bulaşan içermemesi önemlidir. Bu nedenle, bu çalışmada farklı coğrafik bölgelerden elde edilen polen ve arı ekmeği örneklerinde aflotoksin ve okratoksin analizi yapılarak bulaşan içeriğini belirlemek amaçlanmıştır. **Gereç ve Yöntemler:** Bu çalışmada, arıcılardan elde edilen 18 polen (Kırşehir, Nevşehir, Kayseri, Aydın, Silifke, Niğde, Bursa, Denizli, Isparta, Kütahya, İzmir, Afyon) ve 2 arı ekmeğinin (Kırşehir ve Nevşehir) nem, kuru madde, kül, protein ve mikotoksin içerikleri analiz edilmiştir. **Bulgular:** Numunelerin nem içeriği %13,82 ile %29,10 (ortalama 17,80), kuru madde içeriği 70,90 ile 86,18 (ortalama 82,20), kül içeriği %1,60 ile %3,32 (ortalama 2,11) arasında ve protein içeriği ise %15,06 ve %27,95 (ortalama 20,49) arasında bulunmuştur. Aflatoxin B1, Aflatoxin B2, Aflatoxin G1, Aflatoxin G2 ve toplam Aflatoxin (B1+B2+G1+G2) düzeyleri test edilen örneklerde tespit sınırının (0,60 µg/kg) üzerinde bulunmadı. **Sonuç:** Elde edilen sonuçlara göre ülkemizin farklı bölgelerinden alınan polen ve bee bread örneklerinin mikotoksin analiz değerlerinin Türk Gıda Kodeksi Limitlerine uygun olduğu tespit edilmiştir.

**Anahtar Kelimeler:** Polen; arı ekmeği; kimyasal analiz; aflatoxin; mikotoksin

Pollen grain is produced and released from anthers during pollination as the male gametophyte of flowering plants. One of the most important pollinators is honey bee (*Apis mellifera L.*) for entomophilous plants and wind for anemophilous plants.<sup>1</sup> Pollen is potentially a good dietary supplement for human nutrition. A mixture of flower pollen from different species is agglutinated by nectar and honeybee enzymes (e.g. amylase, catalase) secreted by salivary

glands and pollen-loads are formed, which are recognised as bee pollen.<sup>2</sup> Bee bread (Perga), another important bee product, is a natural product obtained by fermenting bee pollen by mixing it with bee saliva and flower nectar in the honeycomb cells of the hive. Bee pollen and bee bread is considered a functional product, having various nutritional values and various bioactive molecules with important biological effects.<sup>3</sup>

**TO CITE THIS ARTICLE:**

Silici S, Keskin U. Mycotoxin analysis in pollen and bee bread samples: Experimental study. J Tradit Complem Med. 2024;7(3):225-31.

**Correspondence:** Sibel SİLİCİ

Erciyes University Faculty of Agriculture, Department of Agricultural Biotechnology, Kayseri, Türkiye

**E-mail:** silicis@erciyes.edu.tr



Peer review under responsibility of Journal of Traditional Medical Complementary Therapies.

**Received:** 29 Feb 2024

**Received in revised form:** 07 May 2024

**Accepted:** 07 May 2024

**Available online:** 15 May 2024

2630-6425 / Copyright © 2024 by Türkiye Klinikleri. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Pollen grain contains a significant amount of secondary plant metabolites such as different phenolic compounds or carotenoids as part of the plant's defense mechanism and has antioxidant activity.<sup>4,5</sup> However, pollen may contain some contaminants.<sup>4</sup> Due to its ideal water (moisture) content, water activity and pH value, pollen often creates an ideal environment for the growth of different microorganisms (bacteria, mold and yeast). As a result of the presence of mold and yeast, mycotoxin production may occur. Mycotoxins are secondary metabolites of different fungal species that are toxic to vertebrates and can cause some diseases and even death. Various mycotoxins [aflatoxins (AFs) B, G and/or M, ochratoxin A (OTA), patulin] have been defined by the European Commission in different foods (nuts, cereals, dried fruit juices, milk, etc.).<sup>6</sup> Maximum tolerable levels were determined for fumonisin B1 and B2, deoxynivalenol and zearalenone. However, there is no information on bee products such as honey, pollen or bee bread. More than a hundred mycotoxins are known, most of which are produced by some species belonging to one of the three fungal species; *Aspergillus*, *Penicillium* and/or *Fusarium*. According to the available literature, it has been shown that the following mycotoxins can be found in pollen; AFs, ochratoxins, fumonisins, zearalenone, deoxynivalenol and its acetoxy derivative, T-2 toxin (T-2), HT-2 toxin, fusarenone-X, diacetoxycirpenol, nivalenol, neosolaniol, roridin A, verrucarrin A,  $\alpha$ - $\beta$  dehydrocurvularin, fomalactone, 6-(1-propenyl)-3,4,5,6-tetrahydro-5-hydroxy-4H-pyran-2-one, 5-[1-(1-hydroxybut-2-enyl)]-dihydrofuran-2-one and 5-[1-(1-hydroxybut-2-enyl)]-furan-2-one.<sup>7-16</sup>

AFs are the product of metabolism of *A. flavus* and *A. parasiticus* strains as *Aspergillus* spp. and different fungal species belonging to the genus *Aspergillus* spp. The most toxic and dangerous AFs are aflatoxins B1 (AFB1) and B2. Both AFB1 and B2 are carcinogenic to humans and animals and are listed as Group 1 carcinogens according to the International Agency for Research on Cancer.<sup>17</sup> It is important to carry out mycotoxicological analyzes together with the microbiological characterization of the pollen. Mycotoxins are secondary metabolites, and even though they occur at ppm and ppb levels, their preva-

lence and threat to human health has been the subject of research.<sup>18,19</sup>

AFs are toxic to mammals, common and the most effective mold-derived carcinogens.<sup>20</sup> It is stated that AFs cannot be completely eliminated by various processing methods and may pose a risk to human health even at a very small rate.<sup>21</sup> Therefore, in this study, it was aimed to determine the mycotoxin content of pollen and bee bread samples obtained from different geographical regions of Türkiye.

## MATERIAL AND METHODS

Pollen (n=18; Kırşehir, Nevşehir, Kayseri, Aydın, Silifke, Niğde, Bursa, Denizli, Isparta, Kütahya, İzmir, Afyon) and bee bread samples (n=2; Kırşehir and Nevşehir) used in the research were obtained from beekeepers. The collected fresh samples were labeled and stored in the freezer at -20 °C in moisture-proof packages until analysis.

### MOISTURE ANALYSIS

For analysis, 5 g of the ground sample was weighed and put in an oven at 130 °C. After the moisture was evaporated, the remaining material was weighed. Moisture content was calculated according to the formula:  $X=(m-m_0)/m \times 100$  (m; initial weight of sample (g),  $m_0$ ; weight of dried sample (g)).

### ASH ANALYSIS

The sample was burned in the muffle furnace at  $550 \pm 10$  °C and turned into ashes and the residue was weighed and the ash amount was calculated.

### PROTEIN ANALYSIS

Protein analysis of the samples was determined using a fully automatic protein analyzer from Dumas (Elementar Analysis Systeme GmbH, Germany). The nitrogen released as a result of burning the sample with pure oxygen (99.9%) at high temperature (850-950 °C) was measured with the help of thermal conductivity and expressed as % protein by multiplying with the appropriate protein factor.<sup>22</sup> Calculation; It was made according to the formula % crude protein (m/m)=%NxF protein (N conversion factor 6.25).

## AFLATOXIN ANALYSIS

High performance liquid chromatography (HPLC), HPLC Pump (Shimadzu LC-20AD) (Shimadzu Corporation, Kyoto, Japan), RF-10AXL Fluorescence detector (Columbia, MA, USA) and HPLC column (ODS3 -250 mm-5 $\mu$ m- 4.6 mm) were used in the analysis. Certified mixed standard with a total aflatoxin concentration of 2,500 ng/mL was used as aflatoxin B1, B2, G1, G2 standard. Each toxin concentration contained 1,000 ng/mL B1, 250 ng/mL B2, 1,000 ng/mL G1, 250 ng/mL G2. AOAC Official Method 991.31 was used for aflatoxin analysis. For the determination of aflatoxin, the quantitative analysis method was applied after the pre-treatments (extraction, cleaning of the extract, immunoaffinity stage, injection into HPLC).<sup>23,24</sup>

Briefly, 25 g sample was weighed and 5 g NaCl was added, 125 mL of 70% methanol (MeOH) was added and mixed at high speed for 5 minutes. It was filtered through coarse filter paper and Whatman no: 4 filter paper using a glass funnel. 5 mL of the filtrate was taken with a pipette and 20 mL of distilled water was added to it. The column was fixed to the injector, 25 mL of extract and water mixture was transferred to the 20 mL injector with a pipette. Then, washing was done by passing 15 mL of distilled water. The flow rate did not exceed 5 mL/min. 1 mL of MeOH is added and slowly passed through the column. Methanol is collected in a 2.5 mL flask. The process is repeated with 1.5 mL of distilled water and the water is taken into the same flask. The total volume is made up to 2.5 mL. Mix thoroughly by vortexing. It is taken into a 2 mL vial and injected. 100  $\mu$ L is injected from each sample once. HPLC conditions; Wavelength Ex: 360 nm; Em: 430 nm, Temperature 40 °C, Pump Flow Rate: 1 mL/min, Pressure<180 bar, Injection Volume 100  $\mu$ L, Column ODS3 column 250 mm 5  $\mu$ m-4.6 mm. Aflatoxin G2 recovery rate=90.72%, G1 recovery rate=94.04%, B2 recovery rate=92.70%, B1 recovery rate=92.24%, Total Aflatoxin recovery rate was 92%.

## OTA ANALYSIS

Shimadzu brand HPLC device was used in the analysis of ochratoxin according to the method of AOAC Official Method 2000.03, R-BIOPHARM Rice

Ochrarep Application Note, Ref no: A23-P14.V2.<sup>25</sup> According to the HPLC-FLD analysis method used for the determination of ochratoxin, its principle is to capture OTA and take it from the IAK with methanol-acetic acid (98:2), under high pressure with the help of a mobile liquid phase, while passing through the HPLC column at certain wavelengths (Ex: 333 nm, Em: 443 nm) was read with a FLD.

## STATISTICAL ANALYSIS

Values are given as the means $\pm$ standard deviation of five measurements.

## RESULTS

The amount of dry matter, ash amount, ash amount in dry matter, moisture content and protein values in the pollen and bee bread samples used in the research are shown in [Table 1](#). The moisture content of the samples varied between 13.82 and 29.10%. The mean moisture content was found to be 17.80 $\pm$ 3.19%. Dry matter content of pollen samples varied between 70.90 and 86.18 values. The average dry matter content was found to be 82.20 $\pm$ 3.19%. The ash content of the samples varied between 1.60% and 3.32%. The average ash content was found to be 2.11 $\pm$ 0.42%. The mean protein content was found to be 20.49 $\pm$ 3.53%.

As a result of the analyzes, Aflatoxin B1 (AFB1), Aflatoxin B2 (AFB2), Aflatoxin G1 (AFG1), Aflatoxin G2 (AFG2) and total Aflatoxin (B1+B2+G1+G2) and OTA not detected ([Table 1](#)).

## DISCUSSION

Evaluation of the microbiological quality of bee pollen is very important as it is used as a supplement in the human diet. Determining the microbiological quality of pollen is as important as determining its chemical content. Although studies of the analysis of mycotoxins in pollen have mostly started in the last decade, the identification of different microorganisms (bacteria, mold and yeast) in pollen samples started with Gilliam's work in the late 1970s.<sup>26</sup>

Although there are Tolerable Daily Intake values for some mycotoxins, there is no information on recommended limits for mycotoxins in pollen. In

**TABLE 1:** Chemical content and mycotoxin analysis of pollen and bee bread samples.

Sample	Dry matter %	Ash %	Ash in dry matter %	Moisture %	Protein %	Total aflatoxin/ ochratoxin A
Pollen 1	70.90	1.82	2.56	29.10	17.22	ND
Pollen 2	84.61	2.24	2.64	15.39	25.84	ND
Pollen 3	84.58	2.76	3.26	15.42	24.44	ND
Pollen 4	80.54	2.12	2.64	19.46	21.40	ND
Beebread 1	78.08	1.86	2.39	21.92	16.48	ND
Pollen 5	83.52	1.74	2.09	16.48	25.89	ND
Beebread 2	84.71	2.24	2.64	15.29	17.04	ND
Pollen 6	85.79	2.31	2.69	14.21	19.81	ND
Pollen 7	86.18	1.76	2.05	13.82	19.83	ND
Pollen 8	84.86	1.74	2.05	15.14	20.81	ND
Pollen 9	82.36	1.72	2.08	17.64	18.57	ND
Pollen 10	84.54	1.96	2.31	15.46	15.11	ND
Pollen 11	84.83	1.94	2.29	15.17	20.12	ND
Pollen 12	77.10	3.32	4.31	22.90	20.72	ND
Pollen 13	84.23	2.11	2.50	15.77	18.83	ND
Pollen 14	85.21	2.19	2.57	14.79	18.86	ND
Pollen 15	83.39	2.05	2.45	16.61	27.95	ND
Pollen 16	84.02	2.26	2.69	15.98	15.06	ND
Pollen 17	78.83	2.46	3.12	21.17	22.06	ND
Pollen 18	75.80	1.60	2.10	24.20	23.80	ND
$\bar{X}\pm SD$	82.20 $\pm$ 3.19	2.11 $\pm$ 0.42	2.57 $\pm$ 0.56	17.80 $\pm$ 3.19	20.49 $\pm$ 3.53	-
Minimum	70.90	1.60	2.05	13.82	15.06	-
Maximum	86.18	3.32	4.31	29.10	27.95	-

ND: Not detected; SD: Standard deviation.

2008, Campos et al. suggested that in the case of AFB1 formation in pollen, the MPC value should be 2 µg/kg, ie 4.2 µg/kg for total AFs.<sup>27</sup> The Tolerable Weekly Intake for ochratoxins as required by the European Food Safety Authority is 0.12 µg/kg bw.<sup>28</sup> In these regulations, there is no information about the recommended limits for mycotoxins in pollen. In our study, the detection limit was determined as 0.60 µg/kg in pollen and bee bread samples. As a result of the analyzes, AFB1, AFB2, AFG1, AFG2 and total aflatoxin (B1+B2+G1+G2) and OTA could not be detected (Table 1).

In a study on this subject, Xue et al. reported that they did not find ochratoxin contamination in Chinese pollen samples, and that this may be due to the collection time and dry weather conditions.<sup>29</sup> Our results was found to be consistent with this study. González et al. detected *Aspergillus* spp. and *Penicillium* spp. in bee pollen from Spain.<sup>30</sup> Ochratoxin contamination has been reported in bee pollen col-

lected in Slovakia.<sup>14</sup> Medina et al. found contamination in bee pollen added to yeast extract sucrose juice inoculated with *A. ochraceus* spores.<sup>9</sup> In a study conducted in Serbia, Kostić et al. determined that there was AFB1 contamination.<sup>31</sup> In another study conducted in Serbia, the number of fungi, incidence of fungi and AFB1 contamination were determined in bee pollen. Examples include *Acremonium* spp., *Alternaria* spp., *Aspergillus* spp., *Cladosporium* spp., *Epicoccum* spp., *Fusarium* spp., *Mucor* spp., *Penicillium* spp. and *Rhizopus* spp. In their studies, the dominance of fungi of the genus *Aspergillus* spp. and *Alternaria* spp. has been shown.<sup>32</sup> Pitta and Markaki showed in their study that mycelial growth was not observed and AFB1 was not detected in bee pollen samples containing natural microbiota.<sup>33</sup>

In this study, mycotoxin analyzes were performed on eighteen different pollen and two different bee bread samples obtained from different regions in Türkiye, and the risks and reliability of the sam-

ples were determined. Analysis results showed that no aflatoxin or ochratoxin derivatives were found in any of the samples. According to the Turkish Food Codex Contaminants Regulation and other international regulations, there is no set limit for pollen and bee bread yet. However, the maximum limits determined for foodstuffs such as dried fruits, peanuts, oil seeds, nuts, hazelnuts, almonds, pistachios and apricot kernels are between 4-15 µg/kg for total aflatoxin and 3-80 µg/kg for OTA. Therefore, the results obtained are below the Turkish Food Codex limits and are safe.<sup>34</sup>

Aflatoxin and ochratoxin results obtained by Campos et al. suggested that the MPC value in pollen is below the value, indicating that the samples can be easily used as a suitable supplementary food.<sup>27</sup>

In addition, since bee pollen is an important risk factor for the formation of ochratoxin and aflatoxin in the food chain, it is important that the samples do not contain harmful formations in the light of the data obtained as a result of the study.

## CONCLUSION

As a result, pollen quality can be significantly affected by the presence of toxigenic fungi. Mycotoxicological analyzes should be included as a regular control measure along with microbiological testing, as the absence of microbial contamination in pollen has not been shown to exclude the presence of mycotoxins. Since AFs and ochratoxins have been

proven to be carcinogenic substances, their presence in pollen is extremely undesirable. Therefore, it is important to monitor mold and mycotoxin levels in feed/food to avoid adverse health effects. It would be beneficial to use pollen as a food supplement in the current legislation. It should cover the tolerable daily/weekly intake for different mycotoxins as well as the sum of the recommended quality parameters. Further studies on the toxicological effect of mycotoxin combinations are needed to obtain reliable and accurate recommendations for pollen quality control. Conducting and disseminating such studies and analyzes is important for the reliability of bee products produced in our country.

### Source of Finance

*This research was supported by University of Erziyes Department of Scientific Research Project (Project no: FYL-2020-10723)*

### Conflict of Interest

*No conflicts of interest between the authors and / or family members of the scientific and medical committee members or members of the potential conflicts of interest, counseling, expertise, working conditions, share holding and similar situations in any firm.*

### Authorship Contributions

*Idea/Concept: Sibel Silici; Design: Sibel Silici; Control/Supervision: Sibel Silici; Data Collection and/or Processing: Uğur Keskin; Analysis and/or Interpretation: Uğur Keskin; Literature Review: Uğur Keskin; Writing the Article: Sibel Silici; Critical Review: Sibel Silici; References and Fundings: Sibel Silici; Materials: Sibel Silici.*

## REFERENCES

- Borg M, Brownfield L, Twell D. Male gametophyte development: a molecular perspective. *J Exp Bot.* 2009;60(5):1465-78. [[Crossref](#)] [[PubMed](#)]
- Campos MGR, Bogdanov S, Bicudo de Almeida-Muradian L, Szczesna T, Mancebo Y, Frigerio CH, et al. Pollen composition and standardisation of analytical methods. *J Apic Res.* 2008;47(2):156-63. [[Crossref](#)]
- Bakour M, Laaroussi H, Ousaaid D, El Ghouizi A, Es-Safi I, Mechchate H, et al. Bee Bread as a Promising source of bioactive molecules and functional properties: an up-to-date review. *Antibiotics (Basel).* 2022;11(2):203. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- Almedia-Muradian LB, Pamiona LC, Coimbra S, Ortrud MB. Chemical composition and botanical evaluation of dried bee pollen pellets. *Journal of Food Composition and Analysis.* 2005;18(1):105-11. [[Crossref](#)]
- Mărgăoan R, Mărghițaș LA, Dezmirean DS, Dulf FV, Bunea A, Socaci SA, et al. Predominant and secondary pollen botanical origins influence the carotenoid and fatty acid profile in fresh honeybee-collected pollen. *J Agric Food Chem.* 2014;62(27):6306-16. [[Crossref](#)] [[PubMed](#)]
- Bennett JW, Klich M. Mycotoxins. *Clin Microbiol Rev.* 2003;16(3):497-516. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- EC Commission. Setting of maximum levels for certain contaminants in foodstuffs-Regulation no: 1881/2006. Official Journal of the EU. 2006; L364, 5-24. Erişim tarihi: 20.11.2024. [[Link](#)]
- Serra-Bonvehí J, Escola Jorda R. Nutrient composition and microbiological quality of honey bee-collected pollen in Spain. *J Agric Food Chem.* 1997;45(3):725-32. [[Crossref](#)]
- Medina A, González G, Saez JM, Mateo R, Jiménez M. Bee pollen, a substrate that stimulates ochratoxin a production by *Aspergillus ochraceus* Wilh. *Systematic and Applied Microbiology.* 2004;27:261-7. [[Crossref](#)] [[PubMed](#)]
- García-Villanova RJ, Cordon C, González Paramás AM, Aparicio P, García Rosales ME. Simultaneous immunoaffinity column cleanup and HPLC analysis of aflatoxins and ochratoxin a in Spanish bee pollen. *J Agric Food Chem.* 2004;52(24):7235-9. [[Crossref](#)] [[PubMed](#)]
- Carrera MA, Miguel E, Fernandez-Alba A, Hernando MD. First survey on the presence of mycotoxins in commercial bee pollen sourced from 28 countries. *Food Control* 2023;152(3):109816. [[Crossref](#)]
- Zaijun L, Zhongyun W, Xiulan S, Yinjun F, Peipei C. A sensitive and highly stable electrochemical impedance immunosensor based on the formation of silica gel-ionic liquid biocompatible film on the glassy carbon electrode for the determination of aflatoxin B1 in bee pollen. *Talanta.* 2010;80(5):1632-7. [[Crossref](#)] [[PubMed](#)]
- Sinkeviciene J, Taraseviciene Z, Tamutis, V. Fusarium Fungi and Mycotoxins in Bee Pollen Collected in Lithuania. *Appl. Sci.* 2023;13(3):1571. [[Crossref](#)]
- Kačániová M, Juráček M, Chlebo R, Kňazovická V, Kadasi-Horáková M, Kunová S, et al. Mycobiota and mycotoxins in bee pollen collected from different areas of Slovakia. *J Environ Sci Health B.* 2011;46(7):623-9. [[Crossref](#)] [[PubMed](#)]
- Cirigliano AM, Rodríguez MA, Godeas AM, Cabrera GM. Mycotoxins from beehive pollen mycoflora. *Journal of Scientific Research and Reports.* 2014;3(7):966-72. [[Crossref](#)]
- Deveza MV, Keller KM, Lorenzon MC, Nunes LM, Sales ÉO, Barth OM. Mycotoxicological and palynological profiles of commercial brands of dried bee pollen. *Braz J Microbiol.* 2015;46(4):1171-6. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- Vidal A, Mengelers M, Yang S, De Saeger S, De Boevre M. Mycotoxin biomarkers of exposure: a comprehensive review. *Compr Rev Food Sci Food Saf.* 2018;17(5):1127-55. [[Crossref](#)] [[PubMed](#)]
- Seo JA, Yu JH. Toxigenic fungi and mycotoxins. An Z. ed. *Handbook of Industrial Mycology.* 1<sup>st</sup> ed. New York: Marcell Dekker; 2005. p.749. [[Crossref](#)]
- Narayanasamy P. *Postharvest Pathogens and Disease Management.* Hoboken, New Jersey: Wiley-Interscience; 2006. p.582. [[Crossref](#)] [[PMC](#)]
- Ellis WO, Smith JP, Simpson BK, Oldham JH. Aflatoxins in food: occurrence, biosynthesis, effects on organisms, detection, and methods of control. *Crit Rev Food Sci Nutr.* 1991;30(4):403-39. [[Crossref](#)] [[PubMed](#)]
- Bullerman LB. Significance of Mycotoxins to Food Safety and Human Health 1,2. *J Food Prot.* 1979;42(1):65-86. [[Crossref](#)] [[PubMed](#)]
- Anderson S. Determination of fat, moisture, and protein in meat and meat products by using the FOSS FoodScan Near-Infrared Spectrophotometer with FOSS Artificial Neural Network Calibration Model and Associated Database: collaborative study. *J AOAC Int.* 2007;90(4):1073-83. [[Crossref](#)] [[PubMed](#)]
- Weaver CM, Trucksess MW. Determination of aflatoxins in botanical roots by a modification of AOAC Official Method 991.31: single-laboratory validation. *J AOAC Int.* 2010;93(1):184-9. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- Şengül Ü. Comparing determination methods of detection and quantification limits for aflatoxin analysis in hazelnut. *J Food Drug Anal.* 2016;24(1):56-62. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- AOAC. Dumas combustion method, AOAC method 992.15. Official Method Of Analysis. 17th ed. 2002. Erişim tarihi: 20.11.2014. Erişim linki: [[Link](#)]
- Gilliam M. Microbiology of pollen and bee bread: the yeasts. *Apidologie.* 1979;10(1):43-53. [[Crossref](#)]
- Campos GRM, Frigerio C, Lopes J, Bogdanov S. What is the future of bee-pollen? *J Apicrop Apimed Sci.* 2010;2(4):131-44. [[Crossref](#)]
- European Food Safety Authority (EFSA). Opinion of the Scientific Panel on contaminants in the food chain of the EFSA on a request from the Commission related to ochratoxin A in food. *The EFSA Journal* 2006;4(6):365. [[Crossref](#)]
- Xue X, Selvaraj JN, Zhao L, Dong H, Liu F, Liu Y. Simultaneous determination of aflatoxins and ochratoxin a in bee pollen by low-temperature fat precipitation and immunoaffinity column clean up coupled with LC-MS/MS. *Food Anal Met.* 2014;7(3):690-6. [[Crossref](#)]
- González G, Hinojo MJ, Mateo R, Medina A, Jiménez M. Occurrence of mycotoxin producing fungi in bee pollen. *Int J Food Microbiol.* 2005;105(1):1-9. [[Crossref](#)] [[PubMed](#)]

31. Kostić AZ, Petrović TS, Krnjaja VS, Nedić NM, Tešić ZL, Milojković-Opsenica DM. et al. Mold/aflatoxin contamination of honey bee collected pollen from different Serbian regions. *J Apic Res.* 2017;56(1):13-20. [[Crossref](#)]
32. Petrović T, Nedić N, Paunović D, Rajić J, Matović K, Radulović Z. et al. Natural mycobiota and aflatoxin B1 presence in bee pollen collected in Serbia. *Biotechnol Anim Husb.* 2014;30(4):731-74. [[Crossref](#)]
33. Pitta M, Markaki P. Study of aflatoxin B1 production by *Aspergillus parasiticus* in bee pollen of Greek origin. *Mycotoxin Res.* 2010;26(4):229-34. [[Crossref](#)] [[PubMed](#)]
34. Resmî Gazete (5.11.2023, Sayı: 32360) sayılı Türk Gıda Kodeksi Bulaşanlar Yönetmeliği; 2023. Erişim tarihi: 20.11.2024. Erişim linki: [[Link](#)]