

A Case of Knotted Foley Catheter in a 59-Year-Old Intensive Care Unit Patient: Challenges and Management

59 Yaşında Yoğun Bakım Hastasında Düğümlenmiş Foley Kateter: Zorluklar ve Yönetimi

¹ Erhan ERDOĞAN^a, ² Abdullah AYDIN^a, ³ Kemal SARICA^{a,b}

^aSancaktepe Şehit Prof. Dr. İlhan Varank Training and Research Hospital, Clinic of Urology, İstanbul, Türkiye

^bBiruni University Faculty of Medicine, Department of Urology, İstanbul, Türkiye

ABSTRACT Urethral catheterization is widely performed procedure in patients of all age groups for various medical purposes, including the collection of urine samples, clean intermittent catheterization, daily urine output monitoring, relieving urinary retention, and performance of a voiding cystourethrography. Although some minor complications namely pain, bleeding, fever etc. can be encountered after this procedure, one rare complication is the formation of knots in the catheter within the bladder. This can occur when a relatively thin, flexible catheter is advanced too far into the bladder. Highly limited cases reported in the literature typically involve young male children where feeding tubes, placed for diagnostic and therapeutic purposes, became knotted. Here, we present a rare complication involving a knotted Foley catheter in the bladder in a 59-year-old female patient being treated in the intensive care unit. We aimed to discuss the encountered challenges and management strategies in light of the available information in the literature.

Keywords: Foley catheter; knotting; catheter complications

ÖZET Üretral kateterizasyon, idrar örneklerinin toplanması, temiz aralıklı kateterizasyon, günlük idrar çıkışı izleme, idrar retansiyonunu giderme ve işeme sistoüretrografisi gerçekleştirme gibi çeşitli tıbbi amaçlarla tüm yaş gruplarındaki hastalarda yaygın olarak yapılan bir prosedürdür. Bu prosedür sonrasında ağrı, kanama, ateş gibi bazı minör komplikasyonlar görülebilse de nadir bir komplikasyon olan mesanede kateterin düğümlenmesi de meydana gelebilir. Bu durum, nispeten ince ve esnek bir kateterin mesaneye fazla itilmesiyle oluşabilir. Literatürde bildirilen son derece sınırlı vakalar, tanısal ve terapötik amaçlarla yerleştirilen beslenme tüplerinin düğümlendiği erkek çocukları içermektedir. Burada, yoğun bakım ünitesinde tedavi gören 59 yaşındaki bir kadın hastada mesanede düğümlenmiş bir Foley kateteri içeren nadir bir komplikasyonu sunuyoruz. Literatürdeki mevcut bilgiler ışığında karşılaşılan zorlukları ve yönetim stratejilerini tartışmayı amaçladık.

Anahtar Kelimeler: Foley kateter; düğümlenme; kateter komplikasyonu

Foley catheters are widely used in clinical practice for urinary drainage. These catheters are made from highly flexible and soft materials to minimize trauma to the urothelium. While they are generally safe for routine use with limited minor problems, complications such as catheter knotting are extremely

rare but can present significant challenges.¹ Flexible catheters can become knotted within the bladder if advanced too far, potentially causing serious urethral trauma when attempts are made to remove them. We present a female case in which a knotted Foley catheter was detected in the bladder.

TO CITE THIS ARTICLE:

Erdoğan E, Aydın A, Sarica K. A case of knotted foley catheter in a 59-year-old intensive care unit patient: Challenges and management. J Reconstr Urol. 2024;14(2):66-8.

Correspondence: Erhan ERDOĞAN

Sancaktepe Şehit Prof. Dr. İlhan Varank Training and Research Hospital, Clinic of Urology, İstanbul, Türkiye

E-mail: erhandr@hotmail.com



Peer review under responsibility of Journal of Reconstructive Urology.

Received: 11 Jul 2024

Accepted: 23 Aug 2024

Available online: 26 Aug 2024

2587-0483 / Copyright © 2024 by Türkiye Klinikleri. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

CASE REPORT

A 59-year-old female patient, admitted to the intensive care unit (ICU) due to respiratory failure, had a Foley catheter inserted upon admission. A few days later, ICU staff encountered difficulties deflating the catheter balloon for removal, and after a failed attempt to remove the catheter, a urology consultation was requested. Bedside ultrasound was performed, and no obvious abnormalities were detected. The Foley catheter was successfully removed with gentle and continuous traction, and it was observed that the Foley catheter had knotted within the bladder (Figure 1). No complications were observed in the patient following the removal of the knotted catheter.

Patient informed consent was obtained to publish his information. The patient's private information remained confidential with the researchers.

DISCUSSION

Urethral catheterization is widely performed in all age group of cases, either temporarily or permanently, for various medical purposes such as collecting urine samples, clean intermittent catheterization, daily urine output monitoring, relieving urinary retention, and voiding cystourethrography. When performed carefully, the complication rate is quite low.¹ The most common complications following urethral catheterization include infection, urothelial trauma, perforation, and creation of false passages. Knotting of the catheter within the bladder is a very rare complication. This condition typically develops due to the use of thin and flexible feeding tubes. The majority of cases reported in the literature are in the pediatric age group.²⁻⁵ It has been reported to occur more frequently in children under the age of two years.¹ Reports of knotted Foley catheters in the bladder of adults are highly limited in the published literature.⁶⁻⁹ The possible mechanism involves the excessive coiling of a long catheter shaft within the bladder when the bladder volume is reduced, allowing the distal end to pass through an open loop and knot upon applying traction. Smaller diameter and longer length of the catheter may contribute to the increased risk of knotting.¹⁰ Approaches in such cases include gentle and continuous traction, urethral dilation and traction,




FIGURE 1: Postoperative view showing the retrieved Foley catheter.

fluoroscopic manipulation with a guidewire, endoscopic removal, and open or percutaneous cystostomy.^{2,6,9,11-13}

The length of Foley catheters used today are standard and are produced in the same measures for both males and females. This is done to prevent the accidental use of a short catheter in males and inflation of the balloon in the urethra. It is recommended that in male patients, the Foley catheter should be fully inserted, and the balloon should be inflated after urine flow is observed. However, in female patients, due to the shorter urethra, full insertion of the catheter is not necessary. The fact that all reported adult cases are female can be explained by the increased likelihood of the catheter coiling on itself because more of the catheter is inserted into the bladder compared to male patients.

The higher incidence of knotting in the pediatric age group can be attributed to the use of thinner and more flexible catheters, as well as the tendency to overestimate the length of the urethra, especially in male children, leading to further advancement of the catheter into the bladder. The over-insertion of the catheter into the bladder is significant. In newborns, the safe insertion length is recommended to be 6 cm in males and less than 2.5 cm in females.¹⁴

In conclusion, although urethral catheterization is considered to be a simple and routine procedure, it should be performed with great care, and the catheter should be inserted only as far as necessary to prevent possible knot formation. In pediatric patients, the use

of flexible and long feeding tubes should be avoided, and the insertion length of the catheter should be limited according to the patient's age and gender. In adult female patients: Physicians need to remember that the urethra is shorter than in male patients, and the Foley catheter should not be advanced to its full length. Adequate training and information for the medical staff performing the placement, observation, and removal of these catheters can be helpful in the prevention of such a limited complication.

Source of Finance

During this study, no financial or spiritual support was received neither from any pharmaceutical company that has a direct connection with the research subject, nor from a company that pro-

vides or produces medical instruments and materials which may negatively affect the evaluation process of this study.

Conflict of Interest

No conflicts of interest between the authors and / or family members of the scientific and medical committee members or members of the potential conflicts of interest, counseling, expertise, working conditions, share holding and similar situations in any firm.

Authorship Contributions

Idea/Concept: Erhan Erdoğan; **Design:** Erhan Erdoğan; **Control/Supervision:** Erhan Erdoğan; **Data Collection and/or Processing:** Abdullah Aydın; **Analysis and/or Interpretation:** Kemal Sarıca; **Literature Review:** Erhan Erdoğan; **Writing the Article:** Erhan Erdoğan, Kemal Sarıca; **Critical Review:** Kemal Sarıca.

REFERENCES

- Singh VP, Sinha S. Spontaneous knotting of urinary catheters placed with nonindwelling intent: case series and literature review. *Urol Ann.* 2019;11(4):443-6. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- Foster H, Ritchey M, Bloom D. Adventitious knots in urethral catheters: report of 5 cases. *J Urol.* 1992;148(5):1496-8. [[Crossref](#)] [[PubMed](#)]
- Gaisie G, Bender TM. Knotting of urethral catheter within bladder: an unusual complication in cystourethrography. *Urol Radiol.* 1983;5(4):271-2. [[Crossref](#)] [[PubMed](#)]
- Klein EA, Wood DP, Kay R. Retained straight catheter: complication of clean intermittent catheterization. *J Urol.* 1986;135(4):780-1. [[Crossref](#)] [[PubMed](#)]
- Turner TWS. Intravesical catheter knotting: an uncommon complication of urinary catheterization. *Pediatr Emerg Care.* 2004;20(2):115-7. [[Crossref](#)] [[PubMed](#)]
- Ball RA, Horton CE Jr, Mandell JA. Transurethral removal of knotted bladder drainage catheter in a male following bladder neck reconstruction. *Urology.* 1993;41(3):234-6. [[Crossref](#)] [[PubMed](#)]
- Ayyıldız A, Huri E, Nuhoğlu B, Germiyanoglu C. Unexpected complication after cystometry in the hypocompliant urinary bladder: formation of a knot in the double lumen urethral catheter—a case report. *Int Urol Nephrol.* 2006;38(3-4):527-9. [[Crossref](#)] [[PubMed](#)]
- Sighinolfi MC, De Stefani S, Micaii S, Mofferdin A, Beato A, Ferrari N, et al. Knotted urethral catheter in an 80-year-old woman: prevention and management of this unusual complication. *Maturitas.* 2008;60(2):180-1. [[Crossref](#)] [[PubMed](#)]
- Brown SC, Lynn NN. An unusual (knotty) complication of clean intermittent self-catheterization in a patient with an augmented bladder. *BJU Int.* 1999;84(4):539. [[Crossref](#)] [[PubMed](#)]
- Raveenthiran V. Spontaneous knotting of urinary catheters: clinical and experimental observations. *Urol Int.* 2006;77(4):317-21. [[Crossref](#)] [[PubMed](#)]
- Sugar EC, Firlit CF. Knot in urethral catheter due to improper catheterization technique. *Urology.* 1983;22(6):673-4. [[Crossref](#)] [[PubMed](#)]
- Harris VJ, Ramilo J. Guide wire manipulation of knot in a catheter used for cystourethrography. *J Urol.* 1976;116(4):529. [[Crossref](#)] [[PubMed](#)]
- Sambrook AJ, Todd A. Untangling of knotted urethral catheters. *Pediatr Radiol.* 2007;37(4):380-3. [[Crossref](#)] [[PubMed](#)]
- Carlson D, Mowery BD. Standards to prevent complications of urinary catheterization in children: should and should-nots. *J Soc Pediatr Nurs.* 1997;2(1):37-41. [[Crossref](#)] [[PubMed](#)]