

Cardiac Toxicity and Death Caused by Muriatic Acid Ingestion

^{id} Enes FERHATLAR^a, ^{id} Burak DEMİRÇİ^a, ^{id} Abuzer ÇOŞKUN^a

^aUniversity of Health Sciences Bağcılar Training and Research Hospital, Department of Emergency Medicine, İstanbul, Türkiye

ABSTRACT Muriatic acid is a solution of hydrogen chloride in water and is classified as a potent and extremely corrosive substance. It is widely used in household cleaning goods such as toilet bowl cleaners, bathroom tile cleaners, and a variety of cleaning tasks around the home. Regarding adults, oral ingestion is commonly associated with suicide or unintentional incidents. This substance is highly dangerous and can cause both local and systemic consequences in an individual's body. The cardiovascular effects of high doses of chemicals such as hydrochloric acid taken for suicidal purposes are a very complex and serious issue. To prevent needless angiography, it is crucial to recognize that ST segment elevation in those with a higher risk of perforation can be attributed to alternative causes. We aimed to contribute to the literature with this case of muriatic acid ingestion, which caused death due to its cardiac and other system effects.

Keywords: Hydrochloric acid; suicide; cardiotoxicity; ST elevation myocardial infarction

Corrosive ingestion is a significant global public health issue.¹ Prevalent in underdeveloped nations and even observed in developed nations. Annually, there are approximately 5,000 to 15,000 cases of caustic ingestions recorded in the USA.² Corrosive ingestion is a critical medical situation that exhibits several clinical manifestations. The clinical progression of the disease is highly intricate. Ingesting corrosive substances can cause significant damage to the lips, oral cavity, throat, and upper airway.

Muriatic acid is a frequently employed chemical for the purpose of cleaning. It primarily consists of hydrochloric acid. Although muriatic acid and hydrochloric acid share the same chemical formula, they are not identical. While both acids contain HCl, hydrochloric acid is in a state of purity, whereas muriatic acid contains conjugated molecules.³ Physical properties of hydrochloric acid, such as boiling and

melting points, density, and pH, depend on the concentration or molarity of HCl in the aqueous solution. They range from those of water at very low concentrations approaching 0% HCl to values for fuming hydrochloric acid at over 40% HCl.^{4,5} The primary approach to managing cases of hydrochloric acid intake is to closely observe the patient and offer symptomatic care. The typical course of treatment typically entails addressing acidosis, providing pain management, and managing haemolysis. Surgical surgery is recommended for suspected occurrences of perforation.^{6,7}

In this context, we discussed a 42-year-old male patient who attempted suicide and was exposed to hydrochloric acid. We present a case of ST-segment elevation coexistence with a subsequent unfavorable prognosis and death, although no significant pathology was found in the coronary angiography results.

TO CITE THIS ARTICLE:

Ferhatlar E, Demirci B, Coşkun A. Cardiac toxicity and death caused by muriatic acid ingestion. Türkiye Klinikleri J Case Rep. 2024;32(3):53-6.

Correspondence: Burak DEMİRÇİ

University of Health Sciences Bağcılar Training and Research Hospital, Department of Emergency Medicine, İstanbul, Türkiye

E-mail: drburakdemirci@hotmail.com



Peer review under responsibility of Türkiye Klinikleri Journal of Case Reports.

Received: 15 Jan 2024 **Accepted:** 15 Feb 2024 **Available online:** 05 Mar 2024

2147-9291 / Copyright © 2024 by Türkiye Klinikleri. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

CASE REPORT

A 42-year-old male patient presented to our emergency department 30 minutes after orally ingesting muriatic acid. The patient stated that he consumed the entire 600 mL of muriatic acid, which contains hydrochloric acid, with the intention of committing suicide while at his workplace, a supermarket. The patient's speech rate and volume, although expressing a persistent desire for water during the interview, were within the usual range. The individual did not provide any accounts of delusions, obsessions, hallucinations, or illusions. Furthermore, they exhibited remorse for their suicidal behavior. His medical history was unremarkable and he was not taking any medication. During the physical examination, clear evidence of a chemical burn in the oral cavity was observed. Bilateral rales were auscultated in the basal regions of the respiratory sounds. The abdominal region was devoid of discomfort and exhibited no indications of rebound tenderness or guarding. The individual's blood pressure measured 110/80 mmHg, their maximal heart rate was 130 beats per minute, their oxygen saturation level was 98%, their respiratory rate was 28 breaths per minute, and their temperature was 37.4 degrees Celsius. The patient was also given ceftriaxone and pantoprazole, with the former administered as an 80 mg bolus and the latter as an 8 mg/hour continuous infusion. Hydration was also administered.

The laboratory data exhibited a general trend of decline, with some significant findings presented in Table 1. The patient's chest X-ray showed no evidence of mediastinitis or subdiaphragmatic free air, nor increased cardiothoracic ratio consistent with pulmonary congestion. Thorax and abdomen computed tomography (CT) imaging revealed bilateral minimal pleural effusion and ground glass areas in both lung parenchyma. No free air densities were noted, suggesting significant perforation in the mediastinum.

In the gastroduodenoscopy performed after the patient's consultation with the relevant department, it was seen that the entire esophagus and stomach were covered with this chemical material. Areas where the mucosa was destroyed were monitored. When water was given, mucosal clearing was not ob-

served and the bulbous and beyond could not be evaluated due to the risk of perforation (Figure 1B). No additional pathology was detected in the patient's electrocardiography (ECG) taken at the time of admission, other than sinus tachycardia and on repeated ECG, there were ST-segment elevations in leads II, III, aVF accompanied by reciprocal ST-segment depressions in leads DI and aVL which was compatible with inferior myocardial infarction (Figure 1A). He was then quickly taken to emergency coronary angiography. As a result of the patient's angiography, no stenosis or thrombosis was observed in the expected location. Coronary arteries were found to be normal.

The patient experienced sudden abdominal pain that worsened over a period of 9 hours. As the abdomen developed tenderness and guarding, repeated CT scans were conducted. The scans revealed fluid heterogeneity and free air within the abdominal cavity. Following a consultation with a surgeon, emergency surgery was deemed necessary for the patient. Cardiac arrest developed in the patient during transfer to the operating room. The patient, who was resuscitated for 60 minutes, did not respond positively and died.

TABLE 1: Laboratory results of the patient at first arrival, 4th hour and 8th hour.

Laboratory parameter	Arrival	4 th hour	8 th hour
Glucose (mg/dL)	222	237	361
Urea (mg/dL)	13.6	25.2	34.5
Creatinine	1.08	1.59	1.92
Sodium (mEq/L)	135	137	131
Potassium (mEq/L)	4.1	4.66	5.23
Calcium (mg/dL)	8.9	9	8.5
Alanine aminotransferase (U/L)	19	72	91
Aspartate aminotransferase (U/L)	31	138	198
Alkaline phosphatase	81	124	143
Amylase (U/L)	23	526	545
Lipase (U/L)	29	1429.7	1150.5
C-reactive protein (mg/dL)	14.47	3.1	29.68
Troponin (ng/L)	17.05	83.77	2461
White blood cell (103/ μ L)	15.44	29.07	31.01
Platelet (103/ μ L)	149	110	110
Activated partial thromboplastin time (sec)	65	85	110.1
International normalized ratio	1.02	2.22	4.8
pH	7.3	7.02	6.69
Lactate (mmol/L)	3	3.1	3.4




FIGURE 1: A) Electrocardiogram image of the patient, B) Image sections from the patient's endoscopy examination.

The study was conducted in accordance with the Declaration of Helsinki and necessary consent was obtained.

DISCUSSION

The ingestion of hydrochloric acid mostly results in harm to the gastrointestinal tract. It is commonly recognized that diluted acids typically result in minimal harm to the mucosal lining, but concentrated alkalis can cause damage to the esophagus. Conversely, it is established that highly concentrated acids can lead to more severe consequences, such as coagulation necrosis and gastrointestinal necrosis.⁸ Additionally, it can lead to metabolic acidosis, hemolysis, renal failure, and mortality. While cardiovascular symptoms resulting from hydrochloric acid use are quite uncommon, it might potentially lead to a life-threatening condition known as myocardial infarction.⁹

The pathophysiology of cardiovascular damage after drinking hydrochloric acid is unclear and very few cases have been reported.¹⁰ The acid-containing household cleaning product mentioned in the article is highly corrosive and generally causes coagulation necrosis which could lead to perforation in the gastrointestinal system and perforation also developed in this patient. It may also cause metabolic acidosis

and possible mechanism is lactic acid accumulation. Development of acidosis after hydrochloric acid ingestion is considered as a poor prognostic sign. Acidosis causes increased oxygen requirement which could lead to myocardial ischemia. It may also cause decreased threshold for arrhythmias including ventricular fibrillation.¹¹

In the case of Islamoglu et al., ST segment elevation was detected in a patient who drank hydrogen peroxide, and no coronary artery occlusion was detected after coronary angiography, and they reported that the factor causing ST segment elevation could be coronary spasm.¹² In a similar case by San Antonio et al., it was emphasized that the factor causing ST segment elevation in a patient who drank hydrochloric acid was not coronary artery occlusion, but acute myocarditis that developed with direct contact between the necrotic upper gastrointestinal tract and the pericardium.¹³ In our case, similar to other similar cases, ST segment elevation was observed, but no obstruction was detected in coronary angiography.

Possible causes of myocardial infarction might be acidosis-induced proinflammatory and prothrombotic state which resulted in acute thrombosis of the coronary artery or coronary plaque rupture triggered by acidosis or coronary artery spasm induced by aci-

dosis or contribution of all this factors in different proportions. As a result, the patient had ECG findings consistent with myocardial infarction and elevated cardiac markers, and these findings need to be explained. Failure to detect thrombus on angiography does not exclude coronary ischemia due to coronary spasm. Additional investigation is required to elucidate the precise methods by which hydrochloric acid exerts its effects.

Source of Finance

During this study, no financial or spiritual support was received neither from any pharmaceutical company that has a direct con-

nection with the research subject, nor from a company that provides or produces medical instruments and materials which may negatively affect the evaluation process of this study.

Conflict of Interest

No conflicts of interest between the authors and / or family members of the scientific and medical committee members or members of the potential conflicts of interest, counseling, expertise, working conditions, share holding and similar situations in any firm.

Authorship Contributions

All authors contributed equally while this study preparing.

REFERENCES

- Lupa M, Magne J, Guarisco JL, Amedee R. Update on the diagnosis and treatment of caustic ingestion. *Ochsner J.* 2009;9(2):54-9. [[PubMed](#)] [[PMC](#)]
- Ekpe EE, Ette V. Morbidity and mortality of caustic ingestion in rural children: experience in a new cardiothoracic surgery unit in Nigeria. *ISRN Pediatr.* 2012;2012:210632. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- Ganapathy VP, Das RR, Chinnakkannan S, Panda SS. An unusual presentation of hydrochloric acid ingestion: a mystery unraveled. *Pediatr Emerg Care.* 2015;31(3):207-8. [[Crossref](#)] [[PubMed](#)]
- Lide D. *CRC Handbook of Chemistry and Physics.* 81st ed. Boca Raton: CRC Press; 2000.
- Perry R, Green D, Maloney J. *Perry's Chemical Engineers' Handbook.* 6th ed. New York: McGraw-Hill Book Company; 1984.
- Chibishev A, Pereska Z, Chibisheva V, Simonovska N. Corrosive poisonings in adults. *Mater Sociomed.* 2012;24(2):125-30. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- Park KS. Evaluation and management of caustic injuries from ingestion of Acid or alkaline substances. *Clin Endosc.* 2014;47(4):301-7. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- Koschny R, Herceg M, Stremmel W, Eisenbach C. Fatal course of a suicidal intoxication with hydrochloric Acid. *Case Rep Gastroenterol.* 2013;7(1):89-96. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- Sari I, Zengin S, Pehlivan Y, Davutoglu V, Yildirim C. Fatal myocardial infarction after hydrochloric acid ingestion in a suicide attempt. *Am J Emerg Med.* 2008;26(5):634.e5-7. [[Crossref](#)] [[PubMed](#)]
- Hung CH, Chen YJ, Chen YC. Occurrence of ST-segment elevation following hydrochloric acid ingestion in a suicide attempt. *Clin Toxicol (Phila).* 2022;60(4):536-8. [[Crossref](#)] [[PubMed](#)]
- Okonek S, Bierbach H, Atzpodien W. Unexpected metabolic acidosis in severe lye poisoning. *Clin Toxicol.* 1981;18(2):225-30. [[Crossref](#)] [[PubMed](#)]
- Islamoglu Y, Cil H, Atilgan Z, Elbey MA, Tekbas E, Yazici M. Myocardial infarction secondary to unintentional ingestion of hydrogen peroxide. *Cardiol J.* 2012;19(1):86-8. [[Crossref](#)] [[PubMed](#)]
- San Antonio R, Pujol López M, Perea RJ, Sabaté M. ST-segment elevation mimicking myocardial infarction after hydrochloric acid ingestion: Acute caustic myocarditis. *J Cardiovasc Comput Tomogr.* 2017;11(1):70-1. [[Crossref](#)] [[PubMed](#)]