

# Letter to the Editor: In the Shadow of Eclipse: Solar Retinopathy

## Editöre Mektup: Tutulmanın Gölgesinde: Solar Retinopati

 Ceren TÜRKOĞLU<sup>a</sup>,  Elif ARSLAN<sup>b</sup>

<sup>a</sup>Lokman Hekim University Faculty of Medicine, Department of Ophthalmology, Ankara, Türkiye

<sup>b</sup>University of Health Science Ulucanlar Eye Training and Research Hospital, Department of Ophthalmology, Ankara, Türkiye

**Keywords:** Vitamin E; coenzyme Q10; solar activity

**Anahtar Kelimeler:** Vitamin E; koenzim Q10; solar aktivite

Solar maculopathy is a rare, well-defined clinical entity which usually occurs in individuals who look directly at the sun or watch an eclipse without any protection. It was observed that prolonged exposure to bright ambient light might cause damage to a thin, cumulative retina at the level of photoreceptors and retinal pigment epithelium. Initial studies on the damage caused by light on the retina were conducted by Deutschmaann<sup>4</sup> and Widemark<sup>3</sup>.

Solar retinopathy was first described by Bechman et al. by visualizing the hyperreflective area in the fovea with optical coherence tomography (OCT).<sup>2</sup> It has already been revealed that the outer retinal layers are more affected in the acute phase of solar retinopathy. However, researchers have discovered that in the current cases series in Türkiye on November 25, 2022, various pathological findings such as foveal reflective areas, deterioration in the photoreceptor layer, and irregularity in the outer retinal layers have been observed.

The outer membrane of photoreceptor cells contains high amounts of polyunsaturated fatty acids and is extremely sensitive to lipid peroxidation. Vitamin E is an effective antioxidant that delays the intracellular accumulation of lipokine pigments, which contain the end products of lipid peroxidation. On the other hand, CoQ10 supports adenosine triphosphate biosynthesis in mitochondria and is an antioxidant that plays an important role in oxidative metabolism and management.<sup>3</sup>

In the current cases with solar retinopathy (7 cases), specific damages in the retinal pigment epithelium, photoreceptor segment and outer retinal layers were detected using spectral domain-OCT imaging in solar retinopathy cases. After the first visit of the cases, coenzyme-Q and vitamin E complex support was started and followed up to 6 months. It was detected that there was no significant decrease in visual acuity was observed in the patients. Following 6 months, it was observed that there was an

**Correspondence:** Ceren TÜRKOĞLU  
Lokman Hekim University Faculty of Medicine, Department of Ophthalmology, Ankara, Türkiye  
**E-mail:** dr.cerenturkoglu@gmail.com



Peer review under responsibility of Türkiye Klinikleri Journal of Ophthalmology.

**Received:** 04 Sep 2024 **Accepted:** 18 Sep 2024 **Available online:** 23 Sep 2024

2146-9008 / Copyright © 2024 by Türkiye Klinikleri. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

improvement in subjective central vision in each patient. Following 6 months, it was observed that there was an anatomical improvement in spectral domain-OCT level in all cases.

Researchers have conducted that solar retinopathy exposed to solar eclipse was examined and it was observed that the changes were in the outer retinal layers and photoreceptor segments. It is considered that the outer limiting membrane is effective in maintaining the best corrected visual acuity level.

As a result of giving vitamin supplements to solar retinopathy patients with outer retinal defects, no pathological findings were detected except for minimal irregularity in the ellipsoid region on OCT.

#### **Source of Finance**

*During this study, no financial or spiritual support was received neither from any pharmaceutical company that has a direct con-*

*nection with the research subject, nor from a company that provides or produces medical instruments and materials which may negatively affect the evaluation process of this study.*

#### **Conflict of Interest**

*No conflicts of interest between the authors and / or family members of the scientific and medical committee members or members of the potential conflicts of interest, counseling, expertise, working conditions, share holding and similar situations in any firm.*

#### **Authorship Contributions**

**Idea/Concept:** Ceren Türkoğlu, Elif Arslan; **Design:** Ceren Türkoğlu, Elif Arslan; **Control/Supervision:** Ceren Türkoğlu; **Data Collection and/or Processing:** Ceren Türkoğlu, Elif Arslan; **Analysis and/or Interpretation:** Ceren Türkoğlu; **Literature Review:** Ceren Türkoğlu; **Writing the Article:** Ceren Türkoğlu; **Critical Review:** Ceren Türkoğlu; **References and Fundings:** Ceren Türkoğlu, Elif Arslan; **Materials:** Ceren Türkoğlu, Elif Arslan; **Diğer:** Ceren Türkoğlu, Elif Arslan.

## REFERENCES

1. Lucas RS, Harper CA, McCombe MF, Mainster MA, Sliney DH, Zamir E. Optical coherence tomography findings in welder's maculopathy. *Retin Cases Brief Rep.* 2007;1(3):169-71. PMID: 25390786.
2. Garg SJ, Martidis A, Nelson ML, Sivalingam A. Optical coherence tomography of chronic solar retinopathy. *Am J Ophthalmol.* 2004;137(2):351-4. PMID: 14962431.
3. Ekicier Acar S, Sarıcaoğlu MS, Çolak A, Aktaş Z, Sepici Dıncel A. Neuroprotective effects of topical coenzyme Q10 + vitamin E in mechanic optic nerve injury model. *Eur J Ophthalmol.* 2020;30(4):714-22. PMID: 30852912.